

SCSL Digital Collections

Diurnal variations in catches of selected species of ichthyoneuston by the Boothbay neuston net off Charleston, South Carolina

| | |
|---------------|--|
| Item Type | Text |
| Publisher | South Carolina State Library |
| Rights | Copyright status undetermined. For more information contact, South Carolina State Library, 1500 Senate Street, Columbia, South Carolina 29201. |
| Download date | 2024-11-13 22:57:20 |
| Link to Item | http://hdl.handle.net/10827/10553 |

- bow) trout, *Salmo gairdneri*. J. Fish. Res. Board Can. 33:826-829.
- COOPER, J. C., A. T. SCHOLZ, R. M. HORRALL, A. D. HASLER, AND D. M. MADISON.
1976. Experimental confirmation of the olfactory hypothesis with homing, artificially imprinted coho salmon (*Oncorhynchus kisutch*). J. Fish. Res. Board Can. 33:703-710.
- HASLER, A. D., AND W. J. WISBY.
1951. Discrimination of stream odors by fishes and relation to parent stream behavior. Am. Nat. 85:223-238.
- NIEMUTH, W.
1967. A study of migratory lake-run trout in the Brule River, Wisconsin: brown trout. Wis. Dep. Nat. Resour. Fish. Manage. Rep. 12, 80 p.
- SCHOLZ, A. T., J. C. COOPER, D. M. MADISON, R. M. HORRALL, A. D. HASLER, A. E. DIZON, AND R. J. POFF.
1973. Olfactory imprinting in coho salmon: behavioral and electrophysiological evidence. Proc. 16th Conf. Great Lakes Res., p. 143-153.
- SCHOLZ, A. T., R. M. HORRALL, J. C. COOPER, AND A. D. HASLER.
1976. Imprinting to chemical cues: the basis for home stream selection in salmon. Science (Wash., D.C.) 192:1247-1249.
- SCHOLZ, A. T., R. M. HORRALL, J. C. COOPER, A. D. HASLER, D. M. MADISON, R. J. POFF, AND R. I. DALY.
1975. Artificial imprinting of salmon and trout in Lake Michigan. Wis. Dep. Nat. Resour. Fish. Manage. Rep. 80, 46 p.
- STUART, T. A.
1957. The migrations and homing behavior of brown trout (*Salmo trutta* L.). Freshwater Salmon Fish. Res. 18, 27 p.

ALLAN T. SCHOLZ

Laboratory of Limnology
University of Wisconsin
Madison, WI 53706

JON C. COOPER

Laboratory of Limnology, University of Wisconsin
Present address: Texas Instruments, Inc.
Bucanan, NY 10511

ROSS M. HORRALL
ARTHUR D. HASLER

Laboratory of Limnology
University of Wisconsin
Madison, WI 53706

DIURNAL VARIATIONS IN CATCHES OF SELECTED SPECIES OF ICHTHYONEUSTON BY THE BOOTHBAY NEUSTON NET OFF CHARLESTON, SOUTH CAROLINA^{1, 2}

The Boothbay neuston net is becoming a standard gear for collection of ichthyoneuston. Sherman and Lewis (1967) reported using this gear for collection

of lobster larvae. Personnel participating in Cooperative Investigations of the Caribbean and Adjacent Regions (CICAR) activities have prepared a "Plan for Sampling the Early Development Stages of Pelagic Fish during CICAR Operations" which describes the use of the neuston net (FAO³). The Boothbay neuston net, initially adopted as the standard for the Marine Resources Monitoring, Assessment and Prediction Program (MARMAP), consists of a pipe frame 2 m wide by 1 m deep with an 8.5-m long net.⁴ Because little was known concerning the sampling performance of this gear, an experiment was designed to test the operating characteristics of two types of frame (galvanized pipe and aluminum pipe) and two lengths of net (4.9 m and 8.5 m with ratios of mouth to open mesh aperture areas of 1:6 and 1:11, respectively). The nets were of 0.947-mm Nitex⁵ mesh.

The results of the experiment defining the operating characteristics of the two types of frame and two lengths of net were described by Eldridge et al. (1977). The present report will describe mainly diurnal variations in catches of ichthyoneuston during the latter experiment, which was conducted during 9-15 July 1973 utilizing the RV *Dolphin*.

Materials and Methods

The neuston net was towed from a boom extending 3 m from the starboard side of the RV *Dolphin*, and the ship was ordered in an arc of radius 1 n.mi. or less to starboard to keep the net mouth out of the ship's wake. The net was towed so that one-half the height (0.5 m) was in the water.

Towing speeds of 1, 2, and 3 m/s were employed with a total of 48 tows being conducted. Twenty-four daylight tows were made between 1107 and 1627 EST and 24 night tows between 2206 and 0432 EST. After setting (which took an average of 29 s), nets were towed 10 min and then retrieved

¹Contribution No. 74 from the South Carolina Marine Resources Center. This work is the result of research sponsored by the MARMAP Program, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service under Contract No. 4-35137. MARMAP Contribution No. 117.

²Contribution No. 451 from the Southeast Fisheries Center, National Marine Fisheries Service, NOAA, Miami, Fla.

³FAO-UNDP Fisheries Program, Mexico City, 1970. A plan for sampling the eggs and larvae of the fishes of Mexican waters. Unpubl. manuscript.

⁴MARMAP is now using a 0.5 × 1 m neuston net.

⁵Reference to trade names does not imply endorsement by the National Marine Fisheries Service, NOAA.

(average time of retrieval was 32 s). After each tow, the catch was drained on a 0.85-mm mesh sieve and preserved in 5% buffered Formalin. Sorting and identification of ichthyoplankton occurred at the Marine Resources Research Institute (MRRI). Fork lengths were measured in forked tail species; total lengths in all others. Relative volume of water strained was determined by the formula: Relative volume strained = (Speed)(Total tow time)(Average fraction of net in water). The reader should consult Eldridge et al. (1977) for further details concerning material and methods as well as the experimental design.

Results

The 4.9-m net was superior to the 8.5-m net in both ease of handling and minimizing damage to

specimens after capture. There was no significant difference in catching ability of the two nets although the 4.9-m net actually caught more specimens during the experiment (Eldridge et al. 1977). The galvanized pipe frame was superior to the aluminum.

A total of 10,621 specimens of ichthyoneuston were collected. The 20 most abundant taxa made up 85.6% (9,088) of the total number of specimens. The remaining 92 taxa composed 14.4% (1,533) of the total (see Table 1 for most numerous taxa collected).

Analyses of variance and covariance tests revealed that total tow duration, speed, and relative volume strained did not vary significantly between day and night tows (Eldridge et al. 1977). Thus, catches between diurnal periods did not appear biased by the conduct of the experiment.

TABLE 1.—Numbers of individuals of selected ichthyoneuston collected in neuston experiment (+ = significantly more abundant for day or night, or no significant difference in catch between day and night at 5% level of significance).

| Taxon | Total number caught | Number in night catches | Number in day catches | Day | Night | Both | Range total length (mm) | Number of tows present |
|--------------------------------|---------------------|-------------------------|-----------------------|-----|-------|------|-------------------------|------------------------|
| <i>Auxis</i> sp. | 3,576 | 3,573 | 3 | | + | | 2-16 | 26 |
| Exocoetidae | 1,245 | 700 | 545 | | + | | 4-83 | 45 |
| Scombridae | 907 | 906 | 1 | | + | | 4-15 | 8 |
| Gerreidae | 513 | 229 | 284 | | | + | 5-14 | 43 |
| Tetraodontidae | 409 | 15 | 394 | + | | | 4-14 | 29 |
| Mullidae | 348 | 7 | 341 | + | | | 5-21 | 29 |
| <i>Mugil curema</i> | 230 | 77 | 153 | + | | | 6-18 | 40 |
| Priacanthidae | 223 | 222 | 1 | | + | | 3-30 | 25 |
| <i>Coryphaena hippurus</i> | 217 | 188 | 29 | | + | | 9-62 | 34 |
| <i>Caranx crysos</i> | 191 | 67 | 124 | + | | | 7-37 | 42 |
| Gobiidae | 180 | 179 | 1 | | + | | 5-14 | 22 |
| Anguilliformes | 143 | 142 | 1 | | + | | 6-84 | 24 |
| Carangidae | 128 | 125 | 3 | | + | | 3-5 | 21 |
| <i>Psenes maculatus</i> | 125 | 125 | 0 | | + | | 6-49 | 20 |
| Hemiramphidae | 124 | 97 | 27 | | + | | 6-57 | 31 |
| <i>Decapterus punctatus</i> | 118 | 46 | 72 | + | | | 24-47 | 32 |
| <i>Monacanthus setifer</i> | 103 | 11 | 92 | + | | | 9-35 | 30 |
| Scorpaenidae | 102 | 92 | 10 | | + | | 3-11 | 29 |
| Holocentridae | 93 | 93 | 0 | | + | | 3-17 | 21 |
| <i>Caranx</i> sp. | 91 | 87 | 4 | | + | | 4-32 | 24 |
| Synodontidae | 88 | 88 | 0 | | + | | 5-32 | 18 |
| <i>Euthynnus alletteratus</i> | 67 | 67 | 0 | | + | | 3-10 | 18 |
| <i>Monacanthus hispidus</i> | 64 | 3 | 61 | + | | | 14-58 | 22 |
| <i>Opisthonema oglinum</i> | 62 | 55 | 7 | | + | | 5-15 | 20 |
| <i>Istiophorus platypterus</i> | 59 | 26 | 33 | | | + | 3-18 | 26 |
| <i>Decapterus</i> sp. | 54 | 53 | 1 | | + | | 7-11 | 18 |
| <i>Coryphaena equisetis</i> | 54 | 50 | 4 | | + | | 8-18 | 21 |
| <i>Aluterus</i> sp. | 49 | 6 | 43 | + | | | 1-105 | 17 |
| <i>Trachinotus falcatus</i> | 48 | 31 | 17 | | + | | 7-18 | 19 |
| Balistidae | 46 | 21 | 25 | | | + | 3-12 | 23 |
| Pomacentridae | 46 | 29 | 17 | | | + | 5-20 | 28 |
| Labridae | 44 | 43 | 1 | | + | | 5-18 | 18 |
| <i>Scomberomorus cavalla</i> | 41 | 39 | 2 | | + | | 5-10 | 18 |
| Serranidae | 40 | 40 | 0 | | + | | 3-16 | 15 |
| Cynoglossidae | 39 | 39 | 0 | | + | | 5-16 | 16 |
| <i>Kyphosus</i> sp. | 39 | 15 | 24 | | | + | 7-21 | 18 |
| <i>Selar crumenophthalmus</i> | 38 | 18 | 20 | | | + | 6-69 | 15 |
| <i>Bothus</i> sp. | 34 | 33 | 1 | | + | | 3-22 | 16 |
| <i>Canthigaster</i> sp. | 33 | 31 | 2 | | + | | 3-17 | 14 |
| <i>Monacanthus</i> sp. | 33 | 3 | 30 | + | | | 8-30 | 11 |
| <i>Dactylopterus volitans</i> | 30 | 3 | 27 | + | | | 8-30 | 11 |
| <i>Seriola</i> sp. | 26 | 4 | 22 | + | | | 5-18 | 14 |
| <i>Seriola rivoliana</i> | 25 | 0 | 25 | + | | | 14-43 | 9 |
| <i>Caranx hippos</i> | 23 | 21 | 2 | | + | | 6-32 | 14 |
| Syngnathidae | 22 | 19 | 3 | | + | | 7-69 | 15 |
| Apogonidae | 22 | 22 | 0 | | + | | 4-10 | 12 |
| <i>Rachycentron canadum</i> | 19 | 19 | 0 | | + | | 6-13 | 11 |

Partial correlation analyses indicated that catches of flyingfishes, Exocoetidae, and silver driftfish, *Psenes maculatus*, were positively correlated with speed. Catches of the planehead filefish, *Monacanthus hispidus*, pygmy filefish, *M. setifer*, and dolphin, *Coryphaena hippurus*, increased with concentrations of sargassum weed which corresponds to earlier observations by Dooley (1972). Catches of Exocoetidae were negatively correlated with manatee grass (Eldridge et al. 1977).

Chi-square analyses indicated that catches of 41 taxa were significantly affected by changes in diurnal period (Table 1). Catches of 29 were greater at night, whereas collections of 12 were greater during daylight hours. There was no evidence to suggest that catches varied significantly between diurnal periods for six species groups.

Data in Table 1 indicate that specimens of *Auxis* sp., Scombridae, Priacanthidae, Gobiidae, Anguilliformes, Carangidae, *Psenes maculatus*, Holocentridae, *Caranx* sp., Synodontidae, *Euthynnus alletteratus*, *Decapterus* sp., *Coryphaena equisetis*, Labridae, *Scomberomorus cavalla*, Serranidae, Cynoglossidae, *Bothus* sp., *Canthigaster* sp., Apogonidae, and *Rachycentron canadum* could be considered "faculative neuston" (Hempel and Weikert 1972). Specimens of Gerreidae, *Istiophorus platypterus*, Balistidae, Pomacentridae, *Kyphosus* sp., and *Selar crumenophthalmus* appear to be "euneuston" as defined by Hempel and Weikert (1972). Similarly, *Mugil curema*, *Caranx crysos*, and *Decapterus punctatus* appear to be "pseudoneuston."

Mugil cephalus was identified as an euneustonic species by Hempel and Weikert (1972); whereas *M. curema* in our samples appeared to be pseudoneustonic. The difference may be real because different species are involved or simply a sampling artifact. Similarly, although young stages of Exocoetidae were reported as rarely encountered and as concentrating at the surface during daytime by Hempel and Weikert (1972), Exocoetidae were the second most abundant taxa in our samples and were taken mostly at night. The reason for this is unknown, but may be due to differences in location, species sampled, or random error associated with sampling of ichthyoneuston.

Tetraodontidae, puffers, were taken most often during the day and were positively correlated with density of manatee grass.

Results of the neuston gear experiment indicated that 1) the 4.9-m net is the superior net for routine surveys, and 2) choice of sampling hours should take into account variation in catches associated with changes in light conditions.

Acknowledgments

We thank members of the crew and scientific party of RV *Dolphin*, who performed the field work, especially Bruce Stender, Bill Leland, and Oleg Pashuk. Thanks are also due to Howard Powles, Paul Sandifer, and Edwin B. Joseph, who reviewed the manuscript and to Patricia Dupree and Lexa Ford who typed the manuscript.

Literature Cited

- DOOLEY, J. K.
1972. Fishes associated with the pelagic sargassum complex, with a discussion of the sargassum community. *Contrib. Mar. Sci.* 16:1-32.
- ELDRIDGE, P. J., F. H. BERRY, AND M. C. MILLER, III.
1977. Test results of the Boothbay neuston net related to net length, diurnal period, and other variables. *S.C. Mar. Resour. Cent. Tech. Rep.* 18, 22 p.
- HEMPEL, G., AND H. WEIKERT.
1972. The neuston of the subtropical and boreal Northeastern Atlantic Ocean. A review. *Mar. Biol. (Berl.)* 13:70-88.
- SHERMAN, K., AND R. D. LEWIS.
1967. Seasonal occurrence of larval lobsters in coastal waters of central Maine. *Proc. Natl. Shellfish. Assoc.* 57:27-30.
- PETER J. ELDRIDGE
*Marine Resources Research Institute
South Carolina Wildlife and Marine Resources Department
P.O. Box 12559, Charleston, SC 29412*
- FREDERICK H. BERRY
*Southeast Fisheries Center
National Marine Fisheries Service, NOAA
75 Virginia Beach Drive, Miami, FL 33149*
- M. CLINTON MILLER, III
*Department of Biometry
Medical University of South Carolina
Charleston, SC 29401*