

South Carolina Department of Health and Environmental Control

ENVIRONMENTAL AFFAIRS

SHELLFISH MANAGEMENT AREA 10A

2022 ANNUAL UPDATE

**Shellfish Sanitation Section
Environmental Affairs
2600 Bull Street
Columbia, SC 29201**

September 2022



WEB ADDRESS
<http://www.scdhec.gov/FoodSafety/ShellfishMonitoring/>

SHELLFISH MANAGEMENT AREA 10A 2022 ANNUAL UPDATE

[Data Through December 2021]



Prepared By:

Ryan Reed, Regional Shellfish Program Manager
Environmental Affairs - Office of Law Enforcement
1362 McMillan Avenue, Suite 300
Charleston, South Carolina 29405

Reviewer:

Mike Marshall, State Shellfish Program Manager
Environmental Affairs – Office of Law Enforcement
927 Shine Avenue
Myrtle Beach, South Carolina 29577

A handwritten signature in black ink, appearing to read 'Mike Marshall', written in a cursive style.

TABLE OF CONTENTS
Shellfish Management Area 10A Annual Update

Summary	2
Introduction.....	3
Pollution Source Survey	7
Survey Procedures	7
Point Source Pollution	7
A. Municipal and Community Waste Treatment Facilities	7
B. Industrial Waste.....	11
C. Marinas.....	11
D. Radionuclides	12
Non-point Source Pollution	12
A. Urban and Suburban Stormwater Runoff.....	12
B. Agricultural Runoff.....	13
C. Individual Sewage Treatment and Disposal Systems	13
D. Wildlife and Domestic Animals.....	13
E. Boat Traffic	13
F. Hydrographic and Habitat Modification	14
Naturally Occurring Pathogens.....	14
A. Marine Biotoxins.....	14
B. <i>Vibrio parahaemolyticus</i>	14
Hydrographic and Meteorological Characteristics	14
Water Quality Studies	15
Conclusions.....	16
Recommendations.....	17
References.....	19

Figures and Tables

Figures:

(1) Shellfish Growing Area 10A.....	20
-------------------------------------	----

Tables:

(1) Shellfish Water Quality Sampling Stations Description.....	21
(2) Fecal Coliform Bacteriological Data Summary Sheet (<i>January 01, 2019 - December 31, 2021</i>).....	22
(3) Fecal Coliform Historical Trend Sheet	23
(4) Water Quality Sampling Station Data.....	24
(5) Rainfall Data (<i>January 01, 2019 - December 31, 2021</i>).....	25
(6) Pollution Event Closures.....	29
(7) Marina Inventory.....	30

2022 ANNUAL UPDATE
Shellfish Management Area 10A

Data Inclusive Dates:
01/01/19 thru 12/31/21

Classification Change:
 X Yes No

Shoreline Survey Completed: Yes

(I)ncreased/(D)ecreased/(N)one:

Prior Report & Date: 2021 Annual Update

 D Approved
 N Conditionally Approved
 I Restricted
 N Prohibited

SUMMARY

Shellfish Management Area 10A (SFMA 10A) appears to be influenced primarily by nonpoint source stormwater runoff. There are several classification changes recommended to be implemented for the 2022-2023 shellfish harvesting season. Stations 10A-08 (Folly River at SC Hwy 171 Bridge), 10A-15 (Secessionville Creek at private docks), 10A-16 (Clark Sound at Ocean View Flats) and 10A-34A (Lighthouse Creek at Secessionville Creek and Clark Sound) failed to meet an Approved classification and will now be classified as Restricted. Stations 10A-15A (Folly Creek at Secessionville Creek), 10A-22 (Folly River State Shellfish Ground at Folly Island), 10A-33 (Lighthouse Creek at Clark Sound) and 10A-35 (Right Fork of Schooner Creek, middle of docks) will now become boundary stations for the upcoming shellfish harvesting season.

During the past several years major storms have impacted SFMA 10A. In September of 2019, Hurricane Dorian produced 6.92 inches of rainfall during a two-day period. On April 24, 2020, a major storm event in SFMA 10A brought heavy rainfall totaling 4.44 inches of rain within a 24-hour period. The area was closed for shellfish harvesting until samples justified reopening the growing area. On June 15, 2020, another storm event brought heavy rainfall totaling 4.88 inches of rain within a 24-hour period. The open shellfish season was already closed during this time and there are no summer harvesting operations operating during this time within SFMA 10A therefore a precautionary closure was not implemented. On July 8, 2021, Tropical Storm Elsa produced 3.11 inches of rain in a 24-hour period in SFMA 10A. Although this tropical storm brought in heavy precipitation to the Charleston area, in SFMA 10A, the total 24-hour precipitation was below the 4-inch closure threshold and no summer harvesting activity was permitted during 2021-2022 in SFMA 10A.

INTRODUCTION

PURPOSE AND SCOPE

The authority to regulate the harvest, sanitation, processing, and handling of shellfish is granted to the South Carolina Department of Health and Environmental Control by Section 44-1-140 of the Code of Laws of South Carolina, 1976, as amended. The Department promulgated Regulation 61-47, which provides the rules used to implement this authority and outlines the requirements applied in regulating shellfish sanitation in the State. This regulation specifically addresses classification of shellfish harvesting areas and requires that all areas be examined by sanitary and bacteriological surveys and classified into an appropriate shellfish harvesting classification.

The United States Food and Drug Administration (USFDA) uses The National Shellfish Sanitation Program's (NSSP) *Guide for the Control of Molluscan Shellfish* to evaluate state shellfish sanitation programs. The NSSP Model Ordinance requires that a sanitary survey be in place for each growing area prior to its use as a source of shellfish for human consumption and prior to the area's classification as Approved, Conditionally Approved, Restricted, or Conditionally Restricted. Each sanitary survey shall be updated on an annual basis and accurately reflect changes which have occurred within the area. Requirement of the annual reevaluation include, at a minimum, field observations of pollution sources, an analysis of water quality data consisting of the past year's data in combination with appropriate previously collected data, review of reports and effluent samples from pollution sources, and review of performance standards for discharges impacting the growing area. A brief report documenting the findings shall also be provided.

The following criteria consistent with the NSSP Model Ordinance and S.C. Regulation 61-47 are used in establishing shellfish harvesting classifications:

Approved Area - Growing areas shall be classified approved when the sanitary survey concludes that fecal material, pathogenic microorganisms, and poisonous or deleterious substances are not present in concentrations that would render shellfish unsafe for human consumption. Approved classifications shall be determined upon a sanitary survey that includes water samples collected from stations in the designated area adjacent to actual or potential sources of pollution. For waters sampled under adverse pollution conditions, the median fecal coliform Most Probable Number (MPN) or the geometric mean MPN shall not exceed fourteen per one hundred milliliters, nor shall more than ten percent of the samples exceed a fecal coliform MPN of forty-three per one hundred milliliters (per five tube decimal dilution). For waters sampled under a systematic random sampling plan, the geometric mean fecal coliform MPN shall not exceed fourteen per one hundred milliliters, nor shall the estimated ninetieth percentile exceed an MPN of forty three per one hundred milliliters (per five tube decimal dilution). Computation of the estimated ninetieth percentile shall be determined using National Shellfish Sanitation Program Guide for the Control of Molluscan Shellfish methodology.

Conditionally Approved Area - Growing areas may be classified conditionally approved when they are subject to temporary conditions of actual or potential pollution. When such

events are predictable, as in non-point source pollution from rainfall runoff or discharge of a major river, a management plan describing conditions under which harvesting will be allowed shall be adopted by the Department prior to classifying an area as conditionally approved. Where appropriate, the management plan for each conditionally approved area shall include performance standards for sources of controllable pollution (e.g., wastewater treatment and collection systems), evaluation of each source of pollution, and means of rapidly closing and subsequently reopening areas to shellfish harvesting. Memorandums of agreements shall be a part of these management plans where appropriate. Shellfish shall not be directly marketed from a conditionally approved area until conditions for an approved classification have been met for a period of time likely to ensure the shellfish are safe for consumption. Shellstock from conditionally approved areas that have been subjected to temporary conditions of actual or potential pollution may be relayed to approved areas for purification or depuration through controlled purification operations only by special permit issued by the Department.

Restricted Area - Growing areas shall be classified restricted when sanitary survey data show a moderate degree of pollution or the presence of deleterious or poisonous substances to a degree that may cause the water quality to fluctuate unpredictably or at such a frequency that a conditionally approved classification is not feasible. Shellfish may be harvested from areas classified as restricted only for the purposes of relaying or depuration and only by special permit issued by the Department and under Department supervision. The suitability of restricted areas for harvesting of shellstock for relay or depuration purposes may be determined using comparison studies of background tissue samples with post-process tissue samples, as well as other process verification techniques deemed appropriate by the Department. For restricted areas to be utilized as a source of shellstock for depuration, or as source water for depuration, the fecal coliform geometric mean MPN of restricted waters sampled under adverse pollution conditions shall not exceed eighty-eight per one hundred milliliters nor shall more than ten percent of the samples exceed a MPN of two hundred and sixty per one hundred milliliters for a five-tube decimal dilution test. For waters sampled under a systematic random sampling plan, the fecal coliform geometric mean MPN shall not exceed eighty-eight per one hundred milliliters nor shall the estimated ninetieth percentile exceed an MPN of two hundred and sixty (five tube decimal dilution). Computation of the estimated ninetieth percentile shall be obtained using National Shellfish Sanitation Program Guide for the Control of Molluscan Shellfish methodology.

Conditionally Restricted Area - Growing areas may be classified conditionally restricted when they are subject to temporary conditions of actual or potential pollution. When such events are predictable, as in the malfunction of wastewater treatment facilities, non-point source pollution from rainfall runoff, discharge of a major river or potential discharges from dock or harbor facilities that may affect water quality, a management plan describing conditions under which harvesting will be allowed shall be prepared by the Department prior to classifying an area as conditionally restricted. Where appropriate, the management plan for each conditionally restricted area shall include performance standards for sources of controllable pollution, e.g., wastewater treatment and collection systems and an evaluation of each source of pollution, and description of the means of rapidly closing and subsequent reopening areas to shellfish harvesting. Memorandums of agreements shall be a part of these management plans where appropriate. Shellfish may be harvested from areas classified as

conditionally restricted only for the purposes of relaying or depuration and only by permit issued by the Department and under Department supervision. For conditionally restricted areas to be utilized as a source of shellstock for depuration, the fecal coliform geometric mean MPN of conditionally restricted waters sampled under adverse pollution conditions shall not exceed eighty-eight per one hundred milliliters nor shall more than ten percent of the samples exceed a MPN of two hundred and sixty per one hundred milliliters for a five-tube decimal dilution test. For waters sampled under a systematic random sampling plan, the fecal coliform geometric mean MPN shall not exceed eighty-eight per one hundred milliliters nor shall the estimated ninetieth percentile exceed an MPN of two hundred and sixty per one hundred milliliters (five tube decimal dilution). Computation of the estimated ninetieth percentile shall be obtained using National Shellfish Sanitation Program Guide for the Control of Molluscan Shellfish methodology.

Prohibited Area - Growing areas shall be classified prohibited if there is no current sanitary survey report or if the sanitary survey report or monitoring data show unsafe levels of fecal material, pathogenic microorganisms, or poisonous or deleterious substances in the growing area or otherwise indicate that such substances could potentially reach quantities that could render shellfish unfit or unsafe for human consumption.

BACKGROUND INFORMATION

This sanitary survey evaluates the current harvesting classification of shellfish growing waters designated as Shellfish Management Area 10A (SFMA 10A). SFMA 10A consists of approximately 15,191 acres of shellfish growing area habitat located in Charleston County, South Carolina. Area 10A extends in a northeasterly to southwesterly direction from the Charleston Harbor to the Stono River. The major water bodies in SFMA 10A include Folly River, Lighthouse Creek, Schooner Creek and Clark Sound. The area is bordered to the northeast by the geographic boundaries of Clark Sound and Parrot Point Creek. An imaginary line from the confluence of the Folly River and the Stono River, northeastward through King Flats, to a point adjacent to Fludd's Creek, at the northwest corner of Clark Sound, defines the area's western boundary. The southern and eastern boundaries consist of the Atlantic Ocean shoreline of Folly and Morris Islands respectively.

The shellfish industry in South Carolina is based primarily on the harvest of the eastern oyster (*Crassostrea virginica*) and hard clams, which include both the northern clam (*Mercenaria mercenaria*) and several small populations of the southern clam (*Mercenaria campechiensis*). The ribbed mussel (*Geukensia demissa*) is also harvested in South Carolina, primarily gathered on a small scale by the general public for recreational harvest. Areas in South Carolina designated for commercial harvest by the South Carolina Department of Natural Resources (SCDNR) include State shellfish grounds, Culture permits, Mariculture permits and Kings Grant areas. The South Carolina Department of Health and Environmental Control disallows the harvesting of shellfish for direct marketing purposes from restricted waters. Shellfish harvesting from Prohibited waters for human consumption is not allowed.

There are seven (7) State Shellfish Grounds (SSGs) within SFMA 10A: S189, S196, S201, S203, S205, S206 East, and S206 West. There is one (1) Recreational Shellfish Ground (R), R201, and there are no Kings Grants (G) located within the area. There are ten (10) Culture Permits (C) and

three (3) Mariculture Permit (M) leases located throughout SFMA 10A.

The shellfish harvesting season in South Carolina typically extends from October 1 through May 31. The South Carolina Department of Natural Resources (SCDNR) has the authority to alter the shellfish harvesting season for resource management purposes and grant permits for year-round mariculture operations. Additionally, the South Carolina Department of Health and Environmental Control has the authority to prohibit shellfish harvesting when necessary to ensure that shellfish harvested in South Carolina waters are safe for human consumption.

The harvesting classifications of SFMA 10A **prior** to this sanitary survey were as follows:

PROHIBITED

1. Those waters and adjacent marshland between 10A-18 from Schooner Creek to the Charleston Harbor;
2. Those waters and adjacent marshland from Bass Creek to the Charleston Harbor;
3. Those waters and adjacent marshland within Parrot Point Creek from Lighthouse Rd down to Schooner Creek and across to the Charleston Harbor;
4. Those waters within 1,000 feet of the old Backman's Commercial Fisheries Dock;
5. Those waters within 1,000 feet of Sunset Cay Marina;
6. Those waters within 1,210 feet of Mariner's Cay Marina;
7. Those waters within 350 feet of Crosby's Commercial Fisheries Dock.

RESTRICTED

1. Those waters and adjacent marshland from Holland Island Creek to Folly Rd (171).
2. The waters of Clark Sound west of Stations 10A-16 and 10A-34A, including Station 10A-16B to the mainland.
3. Station 10A-32 (Block Island Creek at fork) and adjacent waters and marshlands including Station 10A-29 (Block Island Creek at Flats) and Morris Island;
4. Station 10A-15 (Secessionville Creek at private docks) and adjacent waters and marshlands northeast to Station 10A-34A (Lighthouse Creek at Secessionville Creek and Clark Sound.)

CONDITIONALLY APPROVED

None

APPROVED

All other waters in SFMA 10A.

Station Additions/Deactivations/Modifications: None

POLLUTION SOURCE SURVEY

SURVEY PROCEDURES

Shoreline surveys of Shellfish Management Area 10A are conducted by the South Carolina Department of Health and Environmental Controls, Environmental Affairs, Lowcountry – Charleston Shellfish Sanitation Program staff, by watercraft, vehicle, and on foot, during the survey period and are ongoing. Extensive visual examinations of lands adjacent to the waters of SFMA 10A are conducted to determine potential sources of pollution entering shellfish growing waters.

POINT SOURCE POLLUTION

A. Municipal and Community Waste Treatment Facilities

National Pollutant Discharge Elimination System (NPDES) Permitted Facilities			
Permit #	Facility	Outfalls	Permitted Flow (Gallons Per Day)
SC0021229	Chas. Water Systems –Plum Island WWTP	001 – Ashley River	34,797,801 GPD

There are no permitted wastewater treatment plants (WWTP) within SFMA 10A. The James Island Public Service District (PSD) and Charleston Water Systems collection systems together service nearly all of James Island. Public sewer also serves a small portion of Folly Beach as well. The Plum Island WWTP (SC0021229), operated by Charleston Water Systems, receives wastewater from James Island and Folly Beach; however, Plum Island is located on the Ashley River adjacent to Dill Creek, and discharges treated wastewater into the Charleston Harbor (SFMA 10B). The table below summarizes all instances where WWTP facilities exceeded their allowed permit values for their Discharge Monitoring Reports (DMRs), the value of that violation, and flow value. For the 2019-2021 reporting years for this Annual Update, Plum Island WWTP had three instances of permit violations of their DMRs.

WWTP Discharge Monitoring Report Violations 2019-2021					
Facility	Limit	Violation	Outfall	Report Date	Monthly Avg. Flow (Gallons Per Day)
Chas. Water Systems –Plum Island WWTP	501 Enterococci/100 ml	>2420 Enterococci/100ml	001	January 2019	28 MGD
	501 Enterococci/100 ml	1011 Enterococci/100ml	001	March 2019	21 MGD
	501 Enterococci/100 ml	>2420 Enterococci/100ml	001	June 2019	22 MGD

James Island (James Island PSD and Charleston Water Systems), reported forty-six (46) sanitary

sewer overflows (SSO's) for 2019-2021. Folly Beach had no reported SSO's for 2019-2021.

Available hydrographic information suggests the potential of an impact from pollution sources located outside the growing area. The portion of the area from the Charleston Harbor extending southwest to Schooner Creek appears to be adversely impacted during certain hydrographic conditions by waters originating in the Charleston Harbor (Moore, 1984).

Outfalls from wastewater treatment plants discharging into the harbor are subsequently discharging into northeastern portions of SFMA 10A. Flow calculations have established time and distance of travel and place the effluent plumes within Schooner Bay near Fort Sumter. Due to public health concerns, waters from the confluence of Schooner Creek and Schooner Bay and all associated creeks and marshland to Charleston Harbor will remain Administratively Prohibited.

Sanitary Sewer Overflows – 2019-2021				
James Island/Charleston Water Systems				
Date	Location	Gallons Released	Waterbody Entered	Growing Area
7/8/19	2235 Ashley River Rd near Ashley Crossing Ln – West Ashley	1,000	N/A	10B
8/14/19	2169 Harborview Rd, James Island		Marsh near James Island Creek	10B
11/17/19	Lyttleton Ave – West Ashley	2,400	N/A	10B
11/17/19	South Windermere Rd at William Ackerman Ln- West Ashley	15,000	N/A	10B
11/19/19	Marsh near the intersection of I-526 West and Clements Ferry Rd	3,500	Marsh adjacent to Cooper River	10B
12/23/19	Charleston CPW Plum Island – Manhole 39-247	2,460	N/A	10B
12/23/19	Charleston CPW Plum Island – Manhole X-01	4,050	N/A	10B
12/23/19	Charleston CPW Plum Island – Manhole 39	4,440	N/A	10B
12/24/19	Charleston CPW Plum Island – Manhole 11-01	2,910	N/A	10B
12/24/19	Charleston CPW Plum Island - Manhole L1 04	4,500	N/A	10B
12/24/19	Charleston CPW Plum Island – Manhole CC-22	2,100	N/A	10B
1/12/20	CHARLESTON CPW PLUM ISLAND	3,750	N/A	N/A
4/24/20	CHARLESTON CPW PLUM ISLAND	3,850	STORM WATER POND TO MARSH TO ASHLEY RIVER	10B
5/20/20	CHARLESTON CPW PLUM ISLAND	2,400	MARSH LEADING TO ASHLEY RIVER	10B
8/13/20	CHARLESTON CPW PLUM ISLAND	4,000	N/A	N/A
8/25/20	CHARLESTON CPW PLUM ISLAND	3,340	STORM WATER POND TO MARSH TO ASHLEY RIVER	10B
9/18/20	CHARLESTON CPW PLUM ISLAND	1,200	STORM DRAIN TO MARSH TO ASHLEY RIVER	10B
9/18/20	CHARLESTON CPW PLUM ISLAND	600	STORM WATER POND TO MARSH TO ASHLEY RIVER	10B

9/19/20	CHARLESTON CPW PLUM ISLAND	4,800	STORM DRAIN TO ASHLEY RIVER	10B
9/29/20	CHARLESTON CPW PLUM ISLAND	12,000	STORM WATER POND TO MARSH TO ASHLEY RIVER	10B
10/17/20	CHARLESTON CPW PLUM ISLAND	960	N/A	N/A
11/5/20	CHARLESTON CPW PLUM ISLAND	25,800	WEST ASHLEY TIDAL CREEK TO STONO RIVER	11 & 12B
11/5/20	CHARLESTON CPW PLUM ISLAND	4,000	STORM WATER POND NEAR 454 CESSNA DRIVE (NO OUTFALL)	11 & 12B
1/14/2021	Charleston CPW Plum Island – Manhole FF1- 00	150	Possibly ditch to Long Branch Creek	11
2/1/2021	Charleston CPW Plum Island	2,100	Marsh leading to Wappoo Cut to the Ashley River	10B, 11
5/11/2021	Charleston CPW Plum Island Across from 625 WHITE Oaks Dr West Ashley	1,500	Storm Pipe leading possibly to the marsh (approximately 1 mile away)	11
5/19/2021	Charleston CPW Plum Island 160 Main Road in front of Gilligan's Restaurant, West Ashley	100	Drain to an adjacent marsh	11
5/29/2021	Charleston CPW Plum Island Concord St between Laurens Rd & Gadsdenboro St	660	Maybe to Cooper River	10B
6/12/2021	Charleston CPW Plum Island MANHOLE (V1-294)	27,000	Marsh leading to Wappoo Cut to the Ashley River	10B, 11
6/12/2021	Charleston CPW MANHOLE (V1-198)	18,000	Marsh leading to Wappoo Cut to the Ashley River	10B, 11
6/12/2021	Charleston CPW Plum Island MANHOLE (V3-10)	900	Marsh leading to Wappoo Cut to the Ashley River	10B, 11
6/12/2021	Charleston CPW Plum Island MANHOLE (D-10)	150	Ditch leading to Wappoo Cut to the Ashley River	10B, 11
6/13/2021	Charleston CPW Plum Island MANHOLE (58C)	37,530	Storm water collection system leading to the	10B

			Charleston Harbor	
7/8/2021	Charleston CPW Plum Island MANHOLE (39-20)	22,800	Storm water pond to marsh to Ashley River	10B
7/8/2021	Charleston CPW Plum Island	15,600	Marsh to Ashley River	10B
7/8/2021	Charleston CPW Plum Island	600	Storm drain to marsh to Ashley River	10B
7/8/2021	Charleston CPW Plum Island MANHOLE (39-20)	7,500	Storm ditch to creek to marsh to Ashley River	10B
7/8/2021	Charleston CPW Plum Island	1,800	N/A	10B
7/8/2021	Charleston CPW Plum Island	75	Marsh to Ashley River	10B
7/20/2021	Charleston CPW Plum Island 501 Stinson Dr, West Ashley	500	N/A	11
7/22/2021	Charleston CPW Plum Island MANHOLE (39-185)	500	Storm drain to Marsh to Ashley River	10B
7/31/2021	Charleston CPW Plum Island	7,140	Storm Ditch to creek to Ashley River	10B
8/4/2021	Charleston CPW Plum Island Across from 3050 Ashley Town Center Dr-West Ashley	4,500	Marsh leading to West Ashley tidal creek to Stono River	11
8/18/2021	Charleston CPW Plum Island MANHOLE (39-20)	11,400	Storm water pond to marsh to Ashley River	10B
8/18/2021	Charleston CPW Plum Island MANHOLE (S-09.1)	6,900	Storm system to a ditch to marsh to Ashley River	10B
10/5/2021	Charleston CPW Plum Island Magnolia Rd near 1144 Anita Dr & Magnolia Rd at Paula Dr – West Ashley	4,400	Ditch to marsh to Ashley River (a mile away)	10B
FOLLY BEACH				
N/A	N/A	N/A	N/A	None reported

B. Industrial Waste (Discharges) – There are currently no permitted active industrial waste discharges into the waters of SFMA 10A.

C. Marinas - In 2007, prompted by a SCDHEC Office of Coastal Resource Management (OCRM) marina definition change, SCDHEC Shellfish adopted the following marina

definition. S.C. Regulation 61-47, Shellfish defines *Marina* as any of the following: 1) locked harbor facility; 2) any facility which provides fueling, pump-out, maintenance or repair services (regardless of length); or, 3) any facility which has permanent docking space of 250 linear feet or greater. 4) Any water area with a structure which is used for docking or otherwise mooring vessels and constructed to provide temporary or permanent docking space for more than ten boats. 5) A dry stack facility. SCDHEC is currently in the process of identifying all facilities meeting the new marina definition. Once identified, they will be mapped and adequate closure zones established to protect public health.

Prior to the 2007 definition change, there were two recreational marinas located within SFMA 10A. Mariner's Cay Marina is located on the Folly River adjacent to Folly Road. Mariner's Cay has 87 wet slips, approximately 60% of which are currently occupied, and offers fueling and wastewater pump-out services. Sunset Cay Marina (formerly Folly Marina) is located on the Folly River approximately one mile southwest of Mariner's Cay. Sunset Cay Marina has 118 wet slips, approximately 50% of which are currently occupied, and offers wastewater pump-out services. Additionally, two commercial fisheries facilities were located within SFMA 10A. The owner of Backman's Seafood, which is now out-of-business, still operates a dock located in Backman's creek, north of Bowen's Island. This dock has approximately 150 ft. to 200 ft. of dockage and accommodates between 1 and 2 shrimp trawlers. It has above ground diesel fuel tanks, but the tanks are currently empty. Crosby's Fish and Shrimp Company operates a dock located in Folly Creek adjacent to Folly Road. Crosby's has approximately 300 ft. of dockage and typically accommodates 2 shrimp trawlers and 2 fishing vessels. There is no on-site fuel; a fuel truck accomplishes all boat fueling. Table #7 is included at the end of this report, providing additional detail on SFMA 10A boat docking facilities.

- D. Radionuclides** - Sources of radionuclides have not been identified within SFMA 10A, and radionuclide monitoring has not been conducted. No other sources of poisonous or deleterious substances have been identified within the area.

NONPOINT SOURCE POLLUTION

- A. Urban and Suburban Stormwater Runoff** – Past shoreline surveys conducted in SFMA 10A revealed numerous homes adjacent to shellfish growing areas. Single-family homes continue to be built sporadically along the mainland shores as well as on several marsh islands. Heavy development continues to occur around Folly Creek and Oak Island. Run-off from these locations has the potential to affect shellfish growing waters in the Folly River.

The Army Corps of Engineers has not conducted any dredging projects in SFMA 10A recently. The Folly River entrance occasionally requires maintenance dredging. The dredge material is usually either placed on the ocean side of the southernmost portion of Folly Island for beach renourishment or placed in an offshore spoil site. Morris Island is the only dredge spoil area within SFMA 10A. The Army Corps of Engineers reserves the right to use the spoil area at any time. The State Ports Authority (SPA) conducts maintenance dredging adjacent to their terminals (SFMA 10B). The Drum Island spoil

area is the primary site used by the SPA in the Cooper River. The site is reaching its capacity and will soon be permanently closed. The SPA may then use Morris Island more frequently.

The uplands surrounding the shellfish growing waters of SFMA 10A consist of various soil textures defined by the United States Department of Agriculture (USDA), Soil Conservation Service (1971) utilizing general classifications and descriptions. Although lands within SFMA 10A consist of numerous soil types, the area is generally comprised of Wando-Seabrook soils and occur on flat ridges and lower lying bands. The USDA (1971) further describes these soils as "moderately well drained to excessively drained, nearly level to gently sloping, sandy soils."

- B. Agricultural Runoff** - There are no permitted agricultural facilities located in SFMA 10A. The lack of concentrated agricultural activity near the shoreline of the growing waters minimizes the potential for contamination of shellfish waters from agricultural runoff.
- C. Individual Sewage Treatment and Disposal Systems** - Homes adjacent to shellfish growing waters on James Island and Folly Island are primarily served by sanitary sewer, although some homes in the outlying areas, primarily on the northern end of Folly Island, are serviced by individual septic systems. Each system requires inspection by South Carolina Department of Health and Environmental Control's, Environmental Affairs, Bureau of Environmental Health Services Lowcountry – Charleston, On-site Wastewater Section and approval before final installation.
- D. Wildlife and Domestic Animals** - SFMA 10A supports a large population of domestic animals attributable to numerous private residences along the shores of both James Island and Folly Island. The area also supports a moderate amount of wildlife, primarily various types of marine birds and mammals. The entire growing area has an extensive network of small tidal creeks. This creek system provides a possible conduit for animal fecal coliform bacteria to be transported to adjacent growing waters.

Bird Key - Stono Heritage Preserve is a DNR managed heritage preserve, a sand spit island formed in the mouth of the Folly and Stono River. The preserve provides nesting, roosting and foraging habitat for a variety of sea and shore birds. Beginning in the mid-1980's, thousands of eastern brown pelicans, several species of terns, black skimmers, laughing gulls, two species of herons and other incidental species successfully nested on Bird Key Stono every year. Due to bird nesting activity, this Preserve is closed to public use from March 15 thru October 15. Bird Key is a likely contributor to fecal coliform levels in the area during that time however those levels are likely mitigated by Bird Key's proximity to Stono Inlet and the immediate ocean water tidal flushing ocean inlets provide.

- E. Boat Traffic** - Recreational boat traffic is moderate in the area throughout the year. Commercial fisheries boats, ranging in size from 16 to 50 feet, operate throughout the area. There are heavy clam mariculture interests within this area. Folly River and Folly Creek are used on nearly a daily basis by these permit holders. During the recreational

shrimp- baiting season, typically extending from mid-September through mid-November, recreational traffic is heavy.

- F. Hydrographic and Habitat Modification** - Hydrographic and habitat modification in estuarine areas requires both State and Federal approval. Portions of Folly River require maintenance dredging. The United States Army Corps of Engineers also utilize a designated tract of land adjacent to Lighthouse Creek on Morris Island as a dredge spoil site.

NATURALLY OCCURRING PATHOGENS

- A. Marine Biotoxins** - Bivalve shellfish contamination from marine biotoxins has not been shown to be a human health concern within SFMA 10A. During the winter and spring of 1988, South Carolina experienced an occurrence of "Red Tide", specifically *Ptychodiscus brevis* (*K. brevis*), which affected water quality in SFMA 01. There has been no documented reoccurrences of this organism at levels requiring emergency response in South Carolina waters subsequent to the 1988 event. Due to the vast media coverage of events related to *Pfiesteria piscicida*, the Department participates in a State Task Group on Toxic Algae and operates a toxic algae emergency response team. The Department also has a Marine Biotoxin Contingency Plan in place that must be evaluated and updated annually.
- B. *Vibrio parahaemolyticus*** – Because State water temperatures exceed 81 degrees Fahrenheit (F) during June through September, *Vibrio parahaemolyticus* (Vp) management controls must be implemented during these months. Management controls for permitted Aquaculture facilities are specifically addressed in R.61-47. The season for wild-stock harvest is currently closed from June 1 through September 30. The Department is currently opposed to issuance of special wild-stock harvest permits to Certified Shippers during the closed season. Special permit conditions for maricultured triploid oysters during the vibrio control months must include current R.61-47 and NSSP temperature control requirements to be included in the Certified Shipper's HACCP plan.

HYDROGRAPHIC AND METEOROLOGICAL CHARACTERISTICS

PHYSIOGRAPHY

Shellfish Management Area 10A consists of the waters of the Folly River system and the Clark Sound basin. Connections with the Atlantic Ocean are via Charleston Harbor, Lighthouse Inlet and Stono Inlet. Influence of high salinity ocean water entering the area by way of these inlets provide high flow and a subsequent flushing action which assists in maintaining high water quality. The creeks within the area range from 30 to 500 feet in width and average 3 to 25 feet in depth. The entire area is approximately four miles wide (northwest to southeast) and eleven miles long (southwest to northeast).

Tides in SFMA 10A are semidiurnal, consisting of two low and two high tides occurring each lunar day. Mean tidal ranges in Folly Creek, at the Folly Road Bridge, are 5.5 feet during normal

tides and 7.1 feet during spring tides. Wind direction and intensity, as well as atmospheric pressure, typically cause variations in predicted tidal ranges.

Precipitation in SFMA 10A is heaviest during late summer and early autumn. Tropical storms and hurricanes occasionally produce extremely large amounts of rainfall. During winter months heavy rainfall events are uncommon, yet occasional intense thunderstorms associated with rapidly moving low-pressure systems generate heavy rains. Precipitation rarely occurs in the form of snow or ice. Spring weather patterns may be dynamic with associated thunderstorms and severe weather conditions.

In 2017, the collection of rainfall data has been improved for a more consistent, accurate, and reliable data set that can be accessed directly from a shellfish staff member's computer or phone. With assistance from the National Weather Service's, Southeastern River Forecast Center, the development of the South Carolina Shellfish Rainfall Program was introduced and utilized. This new technology provides shellfish program staff with real-time daily updates for rainfall accumulation in each of the South Carolina shellfish growing management areas, as well as providing critical triggers that alert staff to when rainfall thresholds for closures are exceeded.

In September of 2019, Hurricane Dorian produced 6.92 inches of rainfall during a two-day period. On April 24, 2020, a major storm event in SFMA 10A brought heavy rainfall totaling 4.44 inches of rain within a 24-hour period. The area was closed for shellfish harvesting until samples justified reopening the growing area. On June 15, 2020, another storm event brought heavy rainfall totaling 4.88 inches of rain within a 24-hour period. The open shellfish season was already closed during this time and there are no summer harvesting operations operating during this time within SFMA 10A therefore a precautionary closure was not implemented. On July 8, 2021, Tropical Storm Elsa produced 3.11 inches of rain in SFMA 10A which did not meet the requirements for a rainfall closure event. The 2021 precipitation total recorded for SFMA 10A was 49.40 inches.

Prevailing winds along the central portion of the South Carolina coast are from the south and west during spring and summer and from the north during autumn and winter. Wind speeds are generally less than 15 miles per hour (mph); however, strong weather systems may generate winds in excess of 25 mph. Tropical storms and hurricanes occur occasionally.

Freshwater rivers do not discharge directly into SFMA 10A. Freshwater influence is primarily due to rainfall.

WATER QUALITY STUDIES

DESCRIPTION OF THE PROGRAM

The Department currently utilizes a systematic random sampling (SRS) strategy within SFMA 10A in lieu of sampling under adverse pollution conditions. In order to comply with NSSP guidelines, a minimum of thirty samples are required to be collected and analyzed from each station during the review period. Sampling dates are computer generated prior to the beginning of each quarterly period thereby insuring random selection with respect to tidal stage and weather. Day of week selection criteria is limited to Mondays, Tuesdays and Wednesdays due to

shipping requirements and laboratory manpower constraints. Sample schedules are rarely altered.

During July 1998, an updated shellfish water quality data scheduling and collection procedure was formalized. Samples utilized for classification purposes are limited to those samples collected in accordance with the SRS for a 36-month period beginning January 1 and ending December 31. This allows for a maximum of 36 samples per station, yet provides a six-sample “cushion” (above the NSSP required 30 minimum) for broken sample bottles, lab error, breakdowns, etc. This also allows each annual report’s water quality data to meet the requirements for the NSSP Triennial Review sampling criteria.

Nine hundred and twenty-two (922) routine SRS surface water quality samples (<1.0 ft deep) were collected for bacteriological analyses and classification purposes from twenty-eight (28) active water quality sampling stations in SFMA 10A during the period 01/01/19 through 12/31/21. Twenty (20) special samples were also taken for non-classification purposes, associated with reopening after precautionary closures. Samples were collected in 120 ml amber glass bottles, immediately placed on ice and transported to the South Carolina Department of Health and Environmental Control's, Environmental Affairs, Lowcountry – Charleston Laboratory in North Charleston, South Carolina. An additional 120 ml water sample was included with each shipment as a temperature control. At the laboratory, sample sets exceeding a 30-hour holding time or containing a temperature control in excess of 10 degrees C. were discarded (APHA, 1970).

Surface water temperatures are measured utilizing hand-held, laboratory-quality calibrated centigrade thermometers. Salinity measurements are obtained in the laboratory using an automatic temperature compensated refractometer. Additional field data include ambient air temperature, wind direction, tidal stage and date and time of sampling.

MONITORING RESULTS

No stations exceeded a fecal coliform geometric mean MPN value of 14.

Stations 10A-08, 10A-15, 10A-16, 10A-16B, 10A-29A, 10A-34, and 10A-34A exceeded a fecal coliform MPN estimated ninetieth percentile value of 43.

No stations exceeded a geometric mean MPN value of 88 and no stations exceeded a fecal coliform MPN estimated 90th percentile value of 260.

CONCLUSIONS

During this annual review period, SFMA 10A had several sample sites that will be downgraded to a Restricted classification. Based on the review of the fecal coliform bacteriological data and the pollution source survey for SFMA 10A, SFMA 10A appears to be impacted by both point and nonpoint source pollution.

POINT SOURCE POLLUTION

Available hydrographic information suggests the potential of an impact from pollution sources located outside the growing area. The portion of the area from the Charleston Harbor extending southwest to Schooner Creek appears to be adversely impacted during certain hydrographic conditions by waters originating in the Charleston Harbor (Moore, 1984). Outfalls from wastewater treatment plants discharging into the harbor are subsequently discharging into northeastern portions of SFMA 10A. Flow calculations have established time and distance of travel and place the effluent plumes within Schooner Bay near Fort Sumter. Due to public health concerns, waters from the confluence of Schooner Creek and Schooner Bay and all associated creeks and marshland to the Charleston Harbor will remain Administratively Prohibited.

NONPOINT SOURCE POLLUTION

Nearly all of SFMA 10A upland shoreline has residential development bordering the marsh. Impervious surfaces typically result in increased volumes of stormwater runoff and a more rapid movement of stormwater into adjacent shellfish harvesting waters. Increases in rainfall typically result in increased stormwater runoff, which often results in elevated fecal coliform levels. These factors increase the potential for water quality to be adversely impacted within the immediate area.

The majority of stations within SFMA 10A meet the statistical criteria for an Approved classification. Stormwater runoff continues to be the major source of minimal fecal coliform contamination throughout the area. The moderate populations of domestic and wild animals likely impact water quality in the area. Over land runoff in SFMA 10A appears to be mitigated by ocean water flushing through Charleston Harbor, Lighthouse Inlet and Stono Inlet.

RECOMMENDATIONS

There will be four recommended classification changes for the 2022-2023 shellfish harvesting season within SFMA 10A. The below classification are recommended for SFMA:

PROHIBITED

1. Those waters and adjacent marshland between 10A-18 from Schooner Creek to the Charleston Harbor;
2. Those waters and adjacent marshland from Bass Creek to the Charleston Harbor;
3. Those waters and adjacent marshland within Parrot Point Creek from Lighthouse Rd down to Schooner Creek and across to the Charleston Harbor;
4. Those waters within 1,000 feet of the old Backman's Commercial Fisheries Dock;
5. Those waters within 1,000 feet of Sunset Cay Marina;
6. Those waters within 1,210 feet of Mariner's Cay Marina;
7. Those waters within 350 feet of Crosby's Commercial Fisheries Dock.

RESTRICTED

1. Those waters and adjacent marshlands of the Folly River from the Folly River Boat Ramp (10A-08) to Station 10A-22 (Folly River State Shellfish Ground at Folly Island);
2. Those waters and adjacent marshlands of Clark Sound from Station 10A-15 (Secessionville Creek at Private Docks) to Station 10A-33 (Lighthouse Creek at Clark Sound) to Station 10A-35 Right Fork of Schooner Creek, Middle of Docks);
3. Holland Island Creek from SFMA 11 to Folly Rd in SFMA 10A behind Sol Legare Island;
4. Station 10A-32 (Block Island Creek at fork) and adjacent waters and marshlands including Station 10A-29 (Block Island Creek at Flats) and Morris Island.

CONDITIONALLY APPROVED

None

APPROVED

All other waters in SFMA 10A.

Station Additions/Deactivations/Modifications: None

Analysis of sampling data for SFMA 10A demonstrates the probability of a significant impact from rainfall exceeding 4.00" in a 24-hour period. Therefore, a precautionary closure of SFMA 10A will be implemented following rainfall events of greater than 4.00" in a 24-hour period, as measured by the National Weather Service, Southeastern River Forecast Center. This methodology is associated with the concept of the Probable Maximum Precipitation (PMP). PMP estimates for the coastal United States have been published in a series of hydro-meteorological reports (HMRs) by the National Weather Service (*National Weather Service*). PMP estimates for South Carolina's growing areas are derived from HMRs 51, 52, and 53 (*National Research Council, 1985*).

REFERENCES

American Public Health Association, Inc., *Procedures for the Bacteriologic Examination of Sea Water and Shellfish*, 1970, pp. 28-47, *Recommended Procedures for the Examination of Sea Water and Shellfish*, 4th ed., Library of Congress, Washington, D.C.

National Research Council, 1985, "*Safety of Dams - Flood and Earthquake Criteria*," National Academy Press, Washington DC.

National Shellfish Sanitation Program (NSSP), Guide for the Control of Molluscan Shellfish, 2017 Revision. Model Ordinance. United States Food and Drug Administration.

<https://www.fda.gov/media/117080/download>

National Weather Service, The National Oceanic and Atmospheric Administration, *Precipitation Frequency Atlas of the Western US: NOAA Atlas II* - Superintendent of Documents, US Government Printing Office, Washington DC.

NOAA, National Weather Service data base.

South Carolina Department of Health and Environmental Control (SCDHEC), Bureau of Water, 2017, Regulation 61-47, Shellfish. p.9-12.

<https://www.scdhec.gov/sites/default/files/media/document/R.61-47.pdf>

United States Department of Agriculture, Soil Conservation Service, 1971, *Soil Survey of Charleston County, South Carolina*. In cooperation with South Carolina Agricultural Experiment Station and South Carolina Land Resources Conservation Commission, National Cooperative Soil Survey, Washington, D.C. p. 78.

Moore, E., 1984. *Clark Sound: Water Quality Assessment of a Shellfish Growing Area*, South Carolina Department of Health and Environmental Control, Shellfish and Recreational Waters Division.




TABLE #1
Shellfish Management Area 10A
Water Quality Sampling Stations Description

Station	Description
10A-02	Folly Creek at SC Hwy 171 Bridge
10A-04	Folly Creek at Backman Creek
10A-05	King Flats Creek
10A-06	Folly Creek at northern end of Little Island
10A-07	Folly Creek north of old Folly Marina
10A-08	Folly River at SC Hwy 171 Bridge
10A-09	Folly River at Last Dock North
10A-10A	Robbins Creek at the first bend upstream from Cutoff Reach
10A-11	Rat Island Creek at first creek on left from Lighthouse Creek
10A-13	Lighthouse Creek at Folly Creek
10A-15	Secessionville Creek at Private Docks
10A-15A	Folly Creek at Secessionville Creek
10A-16	Clark Sound at Ocean View Flats
10A-16B.....	Clark Sound, 550 yds East of Station 10A-16A
10A-18	Schooner Creek at Schooner Bay
10A-19	Schooner Creek at fork
10A-22	Folly River State Shellfish Ground at Folly Island
10A-23	Lighthouse Creek at First Sister Creek
10A-24	Cole Creek - Second Bend
10A-29A	Block Island Creek at Flats
10A-30	Rat Island Creek - Second Bend
10A-32	Block Island Creek at fork
10A-33	Lighthouse Creek at Clark Sound
10A-34	Secessionville Creek at First Dock nearest Clark Sound
10A-34A	Lighthouse Creek at Secessionville Creek and Clark Sound
10A-35	Right Fork of Schooner Creek, Middle of Docks
10A-36	Unnamed Creek at Fork near RiverFront Subdivision
10A-37	Folly Creek at Oak Island Creek

(Total Active – 28)

TABLE #2

Shellfish Management Area 10A FECAL COLIFORM BACTERIOLOGICAL DATA SUMMARY From Shellfish Water Quality Sampling Stations Between

January 1, 2019 to December 31, 2021

Station #	2	4	5	6	7	8	9	10A	11	13
SAMPLES	33	32	33	33	32	33	33	33	33	33
GEOMEAN	4.8	4	4.2	4.3	4.4	7.2	5.2	4.6	3.4	3
90TH %ILE	18	15	15	14	14	52	16	19	9	8
WATER QLTY	A	A	A	A	A	R	A	A	A	A
CLASSIFICATION	P	A	A	A	A	P	A	A	A	A

Station #	15	15A	16	16B	18	19	22	23	24	29A
SAMPLES	33	33	33	33	33	33	33	33	33	33
GEOMEAN	9.7	3.5	8.9	12.5	4	7.3	3.9	2.7	4.4	9.7
90TH %ILE	47	11	53	89	11	30	14	6	16	71
WATER QLTY	R	A	R	R	A	A	A	A	A	R
CLASSIFICATION	R	R	R	R	P	P	R	A	A	R

Station #	30	32	33	34	34A	35	36	37
SAMPLES	33	33	33	33	33	33	33	33
GEOMEAN	4.1	5.1	4.4	10.3	9.2	8.6	6.2	4
90TH %ILE	17	27	19	54	55	41	36	14
WATER QLTY	A	A	A	R	R	A	A	A
CLASSIFICATION	A	R	R	R	R	R	R	A

A - Approved **CA** - Conditionally Approved **R** - Restricted
RND - Restricted/No Depuration **P** - Prohibited

Table #3
Fecal Coliform Historical Trend Sheet

Area 10A Stations 90thile Values for Annual Updates Related to Rainfall

Station #	2021	2020	2019	2018	2017	2016	2015	2014	2013	2012	2011
10A-02	18	11	7	7	8	12	10	11	6	6	6
10A-04	15	9	5	7	8	10	10	10	6	6	6
10A-05	15	12	7	6	6	7	7	6	4	6	7
10A-06	14	12	9	10	8	10	9	9	7	6	8
10A-07	14	12	9	10	9	7	7	8	7	5	6
10A-08	52	23	9	8	9	9	9	12	9	11	22
10A-09	16	9	9	8	8	6	6	7	7	9	15
10A-10A	19	9	7	8	9	10	9	8	5	5	6
10A-11	9	6	3	4	5	7	7	8	6	6	6
10A-13	8	8	5	5	4	5	6	6	6	6	9
10A-15	47	26	21	27	30	40	31	26	13	11	13
10A-15A	11	6	5	6	8	9	7	6	5	5	6
10A-16	53	34	27	21	26	50	46	35	15	12	16
10A-16B	89	88	67	43	27	39	36	39	24	24	45
10A-18	11	8	8	11	12	16	15	13	7	7	8
10A-19	30	22	15	12	14	15	17	11	10	7	9
10A-22	14	11	9	9	8	6	6	7	6	9	15
10A-23	6	5	4	4	4	5	5	4	4	5	8
10A-24	16	17	14	14	8	11	10	9	4	4	8
10A-29A	71	52	14	15	12	19	23	26	19	14	20
10A-30	17	12	5	5	5	7	8	8	6	5	9
10A-32	27	16	9	12	13	11	11	12	10	12	15
10A-33	19	13	8	8	7	9	9	9	8	8	9
10A-34	54	44	27	24	22	30	25	27	19	16	19
10A-34A	55	39	27	32	32	32	31	34	26	12	22
10A-35	41	29	19	22	25	31	34	34	21	12	24
10A-36	36	29	17	21	15	23	16	23	15	13	22
10A-37	14	8	7	10	11	13	11	10	7	5	10
Annual Rainfall (in inches)	49.40	60.45	47.47	55.25	60.24	45.49	64.58	56.8	27.9	37.4	50.9
ND = No Data Red = Impaired Water Quality											

TABLE #4

**WATER QUALITY
SAMPLING STATIONS DATA**

Shellfish Management Area 10A

Detailed data for each shellfish monitoring station listed in this report's "Fecal Coliform Bacteriological Data Summary Table" and in other shellfish reports, can be obtained by writing South Carolina's Department of Health and Environmental Control – Freedom of Information Office at the address below.

Freedom of Information
SC Dept. of Health & Environmental Control
2600 Bull Street
Columbia, SC 29201

Any explanation or clarity needed on the report's content can be obtained by contacting the preparer(s), and/or reviewer(s) listed on the cover page.

TABLE #5

RAINFALL DATA

Shellfish Management Area 10A

Source:

2019 – 2021 Data

National Weather Service - Southeastern River Forecast Center

Location: Folly Beach, South Carolina

2019 Annual Rainfall Summary
Source: National Weather Service - Southeastern River Forecast Center
Location: Folly Island, South Carolina

2019	JAN	FEB	MAR	APR	MAY	JUNE	JULY	AUG	SEPT	OCT	NOV	DEC
1			0.02			0.01			1.33		0.18	
2			0.04	0.29			0.05					0.07
3			0.01	0.24			0.04	0.58	0.17			
4	0.02	0.07	0.38		0.01		0.05	0.09				
5	0.32		0.20	0.06	0.03	0.37	0.13	0.28	*5.05	0.04		
6			0.03	0.58		0.21	0.68	0.06	*1.87	0.06	0.19	
7				0.15		1.43	0.06			0.09		0.06
8				0.24		0.30	0.02			0.66	0.21	
9				0.04	0.02	0.55		0.03			0.01	
10				0.50	0.02	0.22	0.14					
11						0.03	0.74	0.05				
12		0.17	0.09	0.17		1.00	0.01	0.04				
13		0.07		0.06	0.30	1.45					0.24	
14	0.09				0.04		0.02	0.04		0.12		2.23
15								1.50			0.47	0.02
16		0.12	0.01				0.07	0.03		2.07	0.36	
17		0.04	0.02		0.01			0.59		0.01	0.57	
18	0.05						0.01	1.34	0.01			0.24
19	0.01	0.01		0.24			0.70	0.06		0.01		
20	0.64			0.61		0.04				1.27		
21		0.20				0.32						
22						0.06				0.03		
23						0.66						2.83
24	0.43						0.96				0.10	2.63
25	0.11	0.07						0.41				
26			0.03									
27										0.12		
28		0.01								0.10		
29								0.54				
30	0.04					0.06				0.12		0.97
31								0.02				
Total	1.71	0.76	0.83	3.18	0.43	6.71	3.68	5.66	8.43	4.70	2.33	9.05
*Days highlighted indicate 4 or more inches of rain in a 24-hour period. Blank fields indicate no rainfall.												
* Sample dates are indicated in blue.						ND = No Data			ANNUAL RAINFALL		47.47	

2020 Annual Rainfall Summary
Source: National Weather Service - Southeastern River Forecast Center
Location: Folly Island, South Carolina

2020	JAN	FEB	MAR	APR	MAY	JUNE	JULY	AUG	SEPT	OCT	NOV	DEC
1		0.43		0.70	0.38	0.02	0.58		0.01		0.03	0.82
2							0.02				0.06	
3			0.41				0.24	0.10				
4	0.11		0.01					1.53		0.03		
5	0.10		1.17			0.33		0.08				0.07
6			1.54				0.03	0.05	0.01	0.01		
7		0.77				0.60	0.73	0.28	0.07	0.02	0.05	
8							1.88	0.15			0.02	0.15
9						0.09	0.60	0.07	1.13		0.13	
10				0.01		0.04	0.02		0.18		0.02	
11						0.29	0.04	0.15		1.63	0.52	
12	0.02		0.09			0.02	0.02	0.04	0.09		0.49	
13	0.05			0.01		1.22		0.01			1.77	
14		0.15		1.27		0.01		0.20				
15	0.01	0.03		0.36		*4.88	0.04	0.75	0.03		0.01	
16			0.02	0.40				0.05	0.01	0.16		0.03
17	0.06	0.19							1.50			0.32
18			0.02						0.64			
19		0.57				0.03		0.09				
20	0.01	0.03		1.72	0.13	0.27		0.16				
21		0.55		0.01	1.20	0.06		1.95				0.25
22								0.19		0.08		
23			0.02	0.07	0.18			0.49		0.02		
24			0.21	*4.44		0.64		0.21				0.05
25	0.21	0.70	0.04			0.86	0.32	1.22	0.06	0.22		0.72
26		0.09	0.01		0.20		0.10	0.29	0.51		0.05	
27	0.14	0.13			1.89		0.02	0.88				
28	0.01				0.18	0.07			0.23			
29							0.05	0.05	0.72		0.12	
30	0.33			0.70	0.23	0.07	0.04		0.23	0.18	0.72	
31					0.38		0.24	0.12				0.01
Total	1.05	3.64	3.54	9.69	4.77	9.50	4.97	9.11	5.42	2.35	3.99	2.42
*Days highlighted indicate 4 or more inches of rain in a 24-hour period. Blank fields indicate no rainfall.												
* Sample dates are indicated in blue.						ND = No Data			ANNUAL RAINFALL		60.45	

2021 Annual Rainfall Summary
Source: National Weather Service - Southeastern River Forecast Center
Location: Folly Island, South Carolina

2021	JAN	FEB	MAR	APR	MAY	JUNE	JULY	AUG	SEPT	OCT	NOV	DEC
1	0.15	0.64		0.86				0.10	0.02			
2	0.03		0.06				0.01	0.29				0.01
3	0.18		1.26			0.01	0.45	0.01		0.15		
4			0.23		0.31	0.12		0.43		0.12		
5		0.01			0.04	0.77		0.01				
6		0.12				0.02		0.25	0.12	0.87	0.33	
7		0.45	0.05					0.07	0.02	0.47	0.87	
8	0.63						3.11	0.04		0.04	0.13	0.10
9	0.01							0.20	0.98	0.05		0.93
10		0.04		0.01		0.20	0.61		1.90	0.02		
11							0.01					
12	0.06	0.03			0.65	0.05					0.04	0.17
13		0.16			0.37	3.71		0.01				
14	0.03	0.22				0.40	0.12					
15		1.18					0.06	0.28	0.01			
16	0.20	0.23	0.04			1.29		0.12	0.01			
17						0.28		0.74				
18							0.01	1.01	0.03			
19		0.33	1.59				0.10	0.19	0.03			
20		0.46				0.16	0.17	0.04	0.09			0.07
21			0.10			2.77	0.57	0.02	2.03			0.18
22	0.22	0.02	0.12					0.78	0.48			0.38
23	0.18					0.49	0.31	0.30	0.12		0.02	
24								0.12				
25				1.70						0.69		
26							0.02			0.10	0.02	
27	0.52		0.12			0.09	0.04					
28	0.52						0.75					
29			0.02			0.36	0.64			0.44		
30					0.11							
31			0.01									0.38
Total	2.73	3.89	3.60	2.57	1.48	10.72	6.98	5.01	5.84	2.95	1.41	2.22
*Days highlighted indicate 4 or more inches of rain in a 24-hour period. Blank fields indicate no rainfall.												
* Sample dates are indicated in blue.						ND = No Data			ANNUAL RAINFALL		49.40	

TABLE #6
Shellfish Management Area 10A
Pollution Event Closures
2019 – 2021

Event	Date(s)	Sample Date(s)	Opening Date	Comments
Hurricane Dorian	09/05/2019- 09/06/2019	N/A	N/A	6.92 inches of rain produced during a 2-day period. Open Shellfish Harvesting Season was closed. No summer harvest in SFMA 10A during this time.
April Storm Event	04/24/2020	5/4/2020	5/5/2020	4.03 inches of rain produced during a 24-hr period. Area reopened after special sampling indicated it was safe to reopen the area.
June Storm Event	06/15/2020	N/A	N/A	4.88 inches of rain produced during the storm event. Open Shellfish Harvesting Season was closed. No summer harvest in SFMA 10A during this time.

TABLE #7
Shellfish Management Area 10A
MARINA INVENTORY

Marina	Total Slips	Pump-out Facility	Fuel Dock
Sunset Cay	118	Yes	No
Mariners Cay	89	Yes	Diesel-Gas
Crosby's Fish & Shrimp Co.	7	No	Fuel Truck
Backman's Seafood	3	No	Diesel